

present study and to Dr. D. S. Datar of Central Salt & Marine Chemicals Research Institute, Bhavnagar, for the facilities provided. Thanks also are due to Prof. T. V. Desikachary for the kind permission accorded to the author to study the herbaria at Madras University.

CENTRAL SALT AND MARINE CHEMICALS P. SREENIVASA RAO
RESEARCH INSTITUTE,
BHAVNAGAR,
April 12, 1971.

REFERENCES

- BOERGESEN, F. (1915-20): The marine algae of the Danish West Indies 2. Rhodophyceae. *Dansk. Bot. Arkiv.* 3: 1-504.
- (1934): Some marine algae from the northern part of the Arabian sea with remarks on their geographical distribution. *Det. Kgl. Danske Videnskav Selsk. Biol. Medd.* 11: 1-72.
- (1950): Some marine algae from Maurities additions to the parts previously published II. *ibid.* 18 (11): 1-45.
- DAWSON, E. Y. (1944): The marine algae of the gulf of California. *Allan Hancock Pacific Exp.* 3 (10): 189-452.
- FRICTSCH, F. E. (1945): The structure and reproduction of the algae, vol. 2. Univ. Press. Cambridge. pp. XIV + 939.
- OKAMURA, K. (1926): On the distribution of marine algae in Japan. *Proc. Third Pan. Pacific Sci. Cong.* 1: 958-963.
- SILVA, P. C. (1957): Comparison of algal floristic patterns in the Pacific with those in the Atlantic and Indian oceans, with special reference to *Codium*. *Proc. 9th Pacific Sci. Cong. Bangkok, 1957*, 4: 201-216.

33. RECORD OF *DRAPARNALDIA ACUTA* KUTZ. FROM GUJARAT


(With a text-figure)

The only record of *Draparnaldia* from India is that of Randhawa (1936). He reported *D. plumosa* from 'N. India'. Srinivasan (1965) has made no mention of the genus in his work. During the course of investigation of algal flora of the region *Draparnaldia acuta* Kutz. has been reported from Ahmedabad and for the first time from the country. The alga is briefly described below. The description conforms with that given by Prescott (1951).

Main axis of the thallus bearing horizontal whorled fascicles of branchlets which are ovate to acuminate in outline; cells of the main axis and branchlets are slightly swollen, 34-36 μ in diameter; cells of the branchlets 8 μ in diameter. Chloroplast $\frac{1}{2}$ the length of the cells.

Found in the swamp behind the University, Ahmedabad, on 16th July, 1970.

The main characteristic of the species is the lateral branches forming



a broad spreading fascicle. The branchlets are acute and main axis of the fascicle does not retain its identity.

BOTANY DEPARTMENT,
UNIVERSITY SCHOOL OF SCIENCES,
AHMEDABAD-9,
May 17, 1971.

B. S. VAIDYA
I. A. PATEL

REFERENCES

- PRESCOTT, G. W. (1951): Algae of Western Great Lakes area, Michigan, p. 946.
- RANDHAWA, M. S. (1936): Occurrence and distribution of fresh water algae of North India. *Proc. Indian Acad. Sci.* 4: 36.
- SRINIVASAN, K. S. (1965): Algalum species ex India Oriundae. *J. Bot. Survey India*, 7: 188.